Adult Sleep & Breathing Questionnaire

Date:							
Patient 's Name:							
Patient's Date of Birth:		Age:					
Male Female	e						
Have you ever had a sleep tes	t administered?	yesno					
If yes - when did you have you	ur last sleep test?						
Have you been diagnosed with Sleep Apnea?yesno							
Do you currently use a CPAP or Sleep Appliance for Sleep Apnea?yesno							
Are you happy with your CPAP or Sleep Appliance?yesno							
If you are not happy - why?							
How often do you get out of k	ed to use the rest	room during the night?					
, 0							
Decree and the state of the late of			Yes	No			
Do you usually wake feeling ti	red and unrested:	?					
Do you habitually snore?							
Have you been diagnosted wi	th Hypertension/F	ligh Blood Pressure?					
Do you often suffer from waking headaches?							
Do you regularly experience daytime drowsiness or fatigue?							
Do you have blocked nasal pa							
Has anyone observed you sto							
Do you ever wake up choking							
Do you grind your teeth while							
s your neck circumference greater than 40 cm/ 15.75" ?							
Is your Body Mass Index (BMI) more than 35?						
BMI Formula	BMI =	(your weight in pound	ls X 703)				
	()	our height in inches X your h	eight in inc	hes			

The Epworth Sleepiness Scale

How likely are you to doze off or fall asleep in the following situations, iin contrast to feeling just tired? This refers to your usual way of life in recent times. Even if you have not done some of these things recently, try to work out how they would have effected you. Use the following scale to choose the most appropriate number for each situation.

0 =	no	chance	of	dozi	ng

- 1 = slight chance of dozing
- 2 = moderate chance of dozing
- 3 = high chance of dozing

SITUATION	CHANCE OF DOZING
Sitting and reading	
Watching TV	
Sitting inactive in public place (like a theater or a meeting)	
As a passenger in a car for an hour without a break	
Lying down to rest in the afternoon when circumstances permit	
Sitting and talking to someone	
Sitting quietly after a lunch without alcohol	
In a car, while stopped for a few minutes in traffic	
TOTAL SCORE	

Analyze Your Score

Interpretation:	
-----------------	--

From 0-7 It is unlikely that you are abnormally sleepy

From 8-9 You have an average amount of daytime sleepiness

From 10-15 You may be excessively sleepy, depending on the situation.

You may want to consider seeking medical attention

From 16-20 You are excessively sleep and should consider seeking

medical attention

Berlin Questionnaire

Sleep Apnea

Height (m) Weight	(kg) Age	Male / Female			
Please choose the correct response to each question.					
Category 1		Category 2			
1. Do you snore? a. Yes b. No c. Don't know If you answered 'yes':		6. How often do you feel tired or fatigued after your sleep? □ a. Almost every day □ b. 3-4 times per week □ c. 1-2 times per week □ d. 1-2 times per month □ e. Rarely or never			
2. You snoring is: a. Slightly louder than bre b. As loud as talking c. Louder than talking	eathing	7. During your waking time, do you feel tired, fatigued or not up to par? □ a. Almost every day □ b. 3-4 times per week □ c. 1-2 times per week □ d. 1-2 times per month □ e. Rarely or never			
3. How often do you snore? □ a. Almost every day □ b. 3-4 times per week □ c. 1-2 times per week □ d. 1-2 times per month □ e. Rarely or never	?	8. Have you ever nodded off or fallen asleep while driving a vehicle? □ a. Yes □ b. No If you answered 'yes':			
4. Has your snoring ever be other people? □ a. Yes □ b. No □ c. Don't know	othered	9. How often does this occur? a. Almost every day b. 3-4 times per week c. 1-2 times per week d. 1-2 times per month e. Rarely or never			
5. Has anyone noticed that during your sleep?	you stop breathing	Category 3			
□ a. Almost every day □ b. 3-4 times per week □ c. 1-2 times per week □ d. 1-2 times per month □ □ e. Rarely or never		10. Do you have high bloodpressure?□ Yes□ No□ Don't know			

Scoring Berlin Questionnaire

The questionnaire consists of 3 categories related to the risk of having sleep apnea. Patients can be classified into High Risk or Low Risk based on their responses to the individual items and their overall scores in the symptom categories.

Categories and Scoring:

Category 1: items 1, 2, 3, 4, and 5;

Item 1: if 'Yes', assign 1 point

Item 2: if 'c' or 'd' is the response, assign 1 point

Item 3: if 'a' or 'b' is the response, assign 1 point

Item 4: if 'a' is the response, assign 1 point

Item 5: if 'a' or 'b' is the response, assign 2 points

Add points. Category 1 is positive if the total score is 2 or more points.

Category 2: items 6, 7, 8 (item 9 should be noted separately).

Item 6: if 'a' or 'b' is the response, assign 1 point

Item 7: if 'a' or 'b' is the response, assign 1 point

Item 8: if 'a' is the response, assign 1 point

Add points. Category 2 is positive if the total score is 2 or more points.

Category 3 is positive if the answer to item 10 is '**Yes**' or if the BMI of the patient is greater than 30kg/m₂.

(BMI is defined as weight (kg) divided by height (m) squared, i.e., kg/m₂).

High Risk: if there are 2 or more categories where the score is positive.

Low Risk: if there is only 1 or no categories where the score is positive.

Additional Question: item 9 should be noted separately.